



Characterization of DDX4 Gene Expression in Human Cases with Non-Obstructive Azoospermia and in Sterile and Fertile Mice

Hossein Azizi ^{1*}, Amirreza NiaziTabar ¹, Atiyeh Mohammadi ¹, Thomas Skutella ²

1- Department of Nanobiotechnology, Faculty of Biotechnology, Amol University of Special Modern Technologies, Amol, Iran

2- Institute for Anatomy and Cell Biology, Medical Faculty, University of Heidelberg, Heidelberg, Germany

Abstract

Background: In mammals, spermatogenesis is the main process for male fertility that is initiated by spermatogonial stem cells (SSCs) proliferation. SSCs are unipotent progenitor cells accountable for transferring the genetic information to the following generation by differentiating to haploid cells during spermatogenesis. DEAD-box helicase 4 (DDX4) is a specific germ cell marker and its expression pattern is localized to, spermatocytes, and spermatids. The expression in the SSCs on the basement membrane of the seminiferous tubules is low.

Methods: Immunohistochemistry (IHC) and Fluidigm reverse transcriptase-polymerase chain reaction (RT-PCR) were used to analyze the expression of DDX4 in testis tissue of fertile and sterile mice and human cases with non-obstructive azoospermia.

Results: Our immunohistochemical findings of fertile and busulfan-treated mice showed expression of DDX4 in the basal and luminal compartment of seminiferous tubules of fertile mice whereas no expression was detected in busulfan-treated mice. The immunohistochemical analysis of two human cases with different levels of non-obstructive azoospermia revealed more luminal DDX4 positive cells.

Conclusion: Our findings indicate that DDX4 might be a valuable germ cell marker for analyzing the pathology of germ cell tumors and infertility as global urological problems.

Keywords: DDX4 protein, Seminiferous tubule, Spermatogonial Stem Cell, Testicles.

To cite this article: Azizi H, NiaziTabar A, Mohammadi A, Skutella T. Characterization of DDX4 Gene Expression in Human Cases with Non-Obstructive Azoospermia and in Sterile and Fertile Mice. *J Reprod Infertil.* 2021;22(2):85-91. <http://dx.doi.org/10.18502/jri.v22i2.5793>.

* Corresponding Author:
Hossein Azizi, Faculty of
Biotechnology, Amol
University of Special
Modern Technologies,
Amol, Iran
E-mail:
h.azizi@ausmt.ac.ir

Received: Jun. 10, 2020
Accepted: Sept. 26, 2020

Introduction

The process of spermatozoa production in mammals is defined as spermatogenesis, which occurs within the germ epithelium of the seminiferous tubules in the testis. It starts 5-7 days after birth in rodents and 10-13 years after birth in men and continues throughout life (1).

The seminiferous tubules consist of a unique microenvironment or niche (2) which is a determinative factor for germ cell development (3) consisting of intratubular Sertoli cells (4), peritubular myoid cells, and interstitial Leydig cells (5).

Spermatogonial stem cells (SSCs) are the unipotent progenitor cells of spermatogenesis which differentiate into spermatozoa and consequently transfer the genetic information to the next generation (6-9).

In rodents like mice, spermatogenesis begins by spermatogonia which can renew themselves or differentiate into Ap spermatogonia that are interconnected via cytoplasmic bridges (10). Therefore, Ap spermatogonia divide mitotically to form 4, 8, 16, and sometimes 32 Aal spermatogonia which give rise to more differentiated spermatogonia.

gonia, namely, intermediate and type B spermatogonia (11). Type B spermatogonia undergo the last mitotic division to generate primary spermatocytes which undergo the first meiotic division to produce secondary spermatocytes. Consequently, the second meiotic division finally leads to haploid spermatids (12).

In comparison to mice, there is little information about SSCs in humans. There are three groups of spermatogonia, *i.e.* Adark, Apale, and type B. Adark spermatogonia are considered as reserve stem cells with dense and stainable chromatin while Apale as self-renewing stem cells have less dense and less stainable chromatin which gives rise to type B spermatogonia (13).

The most important form of male infertility is non-obstructive azoospermia (NOA), which occurs due to lack of spermatozoa in the ejaculate and consequently failure of spermatogenesis (14, 15). While there is no treatment to restore the spermatogenesis in NOA patients, inadequate gonadotropin production can be improved by medical treatment. In the recent decade, isolation of similar cells with SSC potential from other sources of stem cell has been considered in many several different studies (16, 17). The possibility of the production of SSCs or Sertoli cells from other types of stem cells is dependent on the advancements in reproduction research in the future.

According to the number of infertile couples in the world and its daily growth rate, using the scientific tools to increase the knowledge in reproductive systems seems crucial. In the recent decade, several investigations were done to find suitable markers of germ cells to introduce new treatments against infertility.

Octamer-binding transcription factor 4 (Oct-4) is a member of Pit-1, Octamer transcription factor (POU) family as a master regulator for stem cell pluripotency. Oct-4 (About 7 kb) is located on chromosome 6 and encodes two distinct variants known as Oct-4A and Oct-4B. It has been proven that Oct-4 is expressed in spermatogonial stem cells (SSCs) in young mice but not in type B SSC (18). Depending on its level of expression, different functions are considered for Oct-4. In mammals, it has an important role during embryogenesis and in humans, it is required for reprogramming somatic cells. In participating with two other transcription factors, SOX2 and NANOG, it controls the self-renewal and pluripotency of embryonic stem cells (ES cells) in mice and humans.

Also, some researches demonstrate that Oct-4 is exclusively expressed in ES cells (21).

A useful marker that is suggested for SSC self-renewal is GFRA (Glial cell line-derived neurotrophic factor family receptor alpha). It has been reported that at low concentration of GFRA, SSCs undergo differentiation phase but the high expression indicates that SSCs undergo self-renewal (22).

In mammals, a unique germ cell marker which is exclusively expressed in the germ cells is DDX4. For the first time, DDX4 gene was discovered in drosophila as a necessary factor for female development (23). Next studies demonstrated that DDX4. (A member of the DEAD (Asp-Glu-Ala-Asp) box family proteins) gene encodes an ATP-dependent RNA helicase which regulates the translation of at least two mRNAs during germ cell development, nanos and gurken (24). The DDX4 gene has a conserved sequence in invertebrates and vertebrates such as *C. elegans*, planarian, xenopus, zebrafish, frog, mouse, rat, cat, pig, rhesus macaques, goat, chicken, and human. The expression of RNA DDX4 in somatic tissues has only been observed during fetal development. Targeted disruption of exon 9 and 10 in a mouse DDX4 homologue (MVH) has shown infertility in males while females were fertile (25).

In this study, we use fertile and infertile mice as model animals and also infertile men with two levels of infertility (non-obstructive azoospermia) to introduce reliable germ cell marker which could be helpful for future studies for diagnosis of male infertility.

Methods

Study population: Testis samples from obstructive azoospermia human were obtained from the Urology Clinics, Medical Faculty, University of Tübingen. Furthermore 3 pairs of mouse testis (C57 BL/6 strain) were used. The animal experiments in this study were approved by the ethical committee of Amol University of special modern technologies (Ir.ausmt.rec.1398.03.07).

SSC isolation: Testis samples from cases with obstructive azoospermia were obtained from the Urology Clinics, Medical Faculty, University of Tübingen, Germany. Furthermore, 3 pairs of mouse testis (C57BL/6 strain) were used. These samples were enzymatically digested with a solution consisting of Dispase (0.5 mg/ml) (Sigma Aldrich, USA), DNase (0.5 mg/ml) (Sigma Al-

drich, USA), Collagenase (0.5 mg/ml) (Sigma Aldrich, USA), and Hank's Balanced Salt Solution (HBSS) (PAA, USA). Digested testicular cells were grown in germ stem cell (GSC) culture media containing 1% N2 supplement (Invitrogen, USA), StemPro-34, 1% L-glutamine (PAA, USA), 60 ng/ml of progesterone (Sigma Aldrich, USA), 30 ng/ml of estradiol (Sigma Aldrich, USA), 5 µg/ml of bovine serum albumin (BSA) (Sigma Aldrich, USA), 30 µg/ml of pyruvic acid (Sigma Aldrich, USA), 1% MEM vitamins (PAA, USA), 8 ng/ml of GDNF (Sigma Aldrich, USA), 1% non-essential amino acids (PAA, USA), 10 ng/ml of FGF (Sigma Aldrich, USA), 100 U/ml of human LIF (Millipore), 1% penicillin/streptomycin (PAA, Laboratories GmbH, USA), 0.1% β-mercaptoethanol (Invitrogen, USA), 1% ES cell qualified FBS, 20 ng/ml of epidermal growth factor (EGF), 6 mg/ml of D+ glucose (Sigma Aldrich, USA), 1 µl/ml of DL-lactic acid (Sigma Aldrich, USA) and 100 µg/ml of ascorbic acid (Sigma Aldrich, USA) at 37°C in an atmosphere of approximately 5% CO₂.

Immunohistochemistry: After decapsulation of tunica albuginea, testicular tissue was washed with PBS and fixed in 4% paraformaldehyde. In the next step, tissue with thickness of 8-10 µm was sliced with a microtome and placed on hydrophilic plus slides and held until used at room temperature. Throughout immunohistochemical staining with DDX4 (Cat No.: ab13840), slides were washed by xylene and replaced by water slowly with a sequence of decreasing ethanol concentrations. Until staining, antigen retrieval was performed using heat-induced epitope retrieval (HIER) at 95°C for 20 min and non-specific binding sites in the tissue sections were blocked with 10% BSA/0.3% Triton in PBS.

Reverse transcriptase polymerase chain reaction (RT-PCR): The expression level of DDX4 gene in the mice testicular cells was examined by the Fluidigm Biomark system. Testicular germ cells were picked up with a micromanipulator, lysed with a solution of lysis buffer consisting of 2.5 µl 0.2× assay pool, 9 µl of RT-PreAmp Master Mix (Invitrogen, USA), 0.2 µl of SuperScript III RT/Taq (Invitrogen, USA) and 1.3 µl of TE buffer., Next, amplified output of RNA-targeted copies with TaqMan was analyzed by BioMark realtime quantitative PCR (qPCR). Two technical repeat measurements were taken to examine the samples. The normalization of data was carried out using DDX4

(Mm00802445_m1) and mRNA expression fold change compared to MEF feeder cells. Using Excel and GenEx applications, the Ct values were determined.

Results

Immunohistochemical analysis of the testis tissue obtained from two human patients with different levels of infertility (Non-obstructive azoospermia) showed different numbers of DDX4 positive cells with different DDX4 expression levels, one with higher (Figure 1; A1) and the other with lower signal intensity (Figure 1; B1). In both cases, the cells were more condensed in the luminal compartment. Also, comparative immunohistochemistry analysis of seminiferous tubules of mice showed that DDX4 was expressed both in the basal and luminal compartment of seminiferous tubules (Undifferentiated and differentiated part) of fertile mice (Figure 2; A1). In contrast, in busulfan-treated mice, no expression of DDX4 was detected (Figure 2; B2). A normal pattern of expression of DDX4 and an abnormal pattern was observed in fertile (Figure 2; B1) and sterile mice (Figure 2; A1), respectively.

Bright field analysis of testicular cells of infertile human and sterile mouse (Figure 3; A, B) and fertile mouse (Figure 3; C) showed the morphological characteristics of these cells after extraction and expansion. Also, immunohistochemical staining with DDX4 (Figure 3; C3) would be useful for attributing morphological features to positive germ cells.

Immunohistochemical analysis of 1 week old mice revealed luminal expression of DDX4 (Fig-




Figure 1. Immunohistochemical analysis of two abnormal human testicular tissues from infertile men. Low basal and high luminal DDX4 expression in case 1 (Merge, A1) and moderate expression in case 2 in luminal cells (Merge, B1). Dapi=blue nuclear staining (A3, B3)




Figure 2. Immunohistochemical analysis of sterile and fertile mice. Non-detectable expression of DDX4 in busulfan-treated mice (Merge, A1). Expression of DDX4 in the germinal epithelium of fertile mice (Merge, A1)




Figure 3. Bright field images of testicular cells in sterile and fertile mice and infertile human *in vitro*. Testicular cells from sterile mouse after expansion in culture (A). Infertile human testicular cell expansion in culture (B). Immunohistochemical analysis of fertile mouse testicular cells. Bright field of testicular cells of fertile mice (C1), testicular cells of fertile mice stained with blue DAPI (C2), testicular cells of fertile mice stained with DDX4 (C3), and merge of DAPI and DDX4 (C4)

ure 4; A). On the other hand, immunohistochemical analysis of 4 week old mice indicated that the




Figure 4. Immunohistochemical analysis and RT-PCR of DDX in 1 and 4 week old mice. Luminal expression of DDX4 in seminiferous tubules of 1 week old mouse (A). The expression pattern of DDX4 in a 4 week old mouse showed that the DDX4 positive cells were redistributed from the lumen to the basal compartment of seminiferous tubules (B). Significantly increased expression of DDX4 in 4 week old mouse testis in comparison to 1 week old mice by Fluidigm real-time PCR analysis (C)

cells with DDX4 expression were shifting from the luminal to the undifferentiated basal part of the seminiferous tubules, typical for postnatal testis development (Figure 4; B). Also, Fluidigm real-time PCR indicated a significantly higher expression of DDX4 in 4 week old mice (Figure 4; C).

Discussion

Spermatogenesis is responsible for sperm production and consequently transportation of genetic information to the next generation. DDX not only have roles in spermatogenesis but also is crucial for primordial germ cell (PGC) formation which anomalies in differentiate of PGSs could lead to dysfunction of spermatogenesis (26-30). Expression of DDX4 has been shown in various species during spermatogenesis but Wenxu Zhu et al. demonstrated that expression of DDX4 was seen in spermatogonia and spermatocytes while no expression was detected in spermatids and spermatozoas (31).

Although many researchers claimed that expression of DDX4 is restricted to germ cells but M.

Lejong et al. proved that in chick embryo, in addition to germ cells, DDX4 protein was detected in somatic cells, too (32).

In the last 10 years, finding a useful *in vitro* tool has been a main objective of reproductive biology research. MACS (Magnetic-activated cell sorting) and FACS (Fluorescence-activated cell sorting) are two of the common *in vitro* assays used in several investigations. In a recent study, MACS was used to enrich undifferentiated spermatogonia from human testicular cell suspensions (33, 34). Also, in other recent studies, FACS was applied for the characterization of mouse adult testicular macrophage populations (35). In a recently study on gene editing on SSCs mediated by CRISPR-Cas9, c-kit and GFR α were used as characterization marker (36). In this present study, we clearly showed that DDX4 would use as a specific marker for undifferentiated spermatogonia.

Immunohistochemistry results indicated that the expression of DDX4 in undifferentiated seminiferous tubule cells (Spermatogonia) was positive. Similarly, Abbasi et al. reported that in immunohistochemistry analysis of THY1 (A conserved marker of SSCs of bovine, rodents, and primates) positive cells were all positive for DDX4 (37). Several investigations have analyzed DDX4 expression in normal human and mice (24, 38). In the present study, comparison of two different abnormal cases with different levels of tissue disruption was done which showed that the expression of spermatogonia had decreased. Similarly, Lee et al. reported that, in prepubertal canine (1-3 months), DAZL (As an undifferentiated marker of male germ cell) was expressed in luminal cells but at pubertal age (4 months), these cells were located near the basement membrane of the seminiferous tubules as detected by immunohistochemistry analysis (39). Also overexpression of DDX4 was seen in human cancers like ovarian cancer and neuroblastoma (40).

Also, our immunohistochemistry analysis of two human abnormal cases with different levels of expression of DDX4 has proven that this marker would be useful for future clinical treatments. In a recent clinical study, pathological and immunohistochemical analyses were performed with 145 cases with a testicular germ cell tumor (TGCT) (41). In some kinds of infertility in men like oligozoospermia, the expression of DDX4 in declined dramatically (42). Analysis of sperm mRNA like DDX4 mRNA by micro-TESE (micro testicular sperm extraction) is a functional way to predicting

of successful sperm retrieval in nonobstructive azoospermic patients (43). In summary, we claimed that DDX4 could be used as a molecular marker for investigation of spermatogenic disorders.

Conclusion

In this study, DDX4 expression was detected in spermatogonia stem cells in normal, but not in the busulfan-treated mice. Furthermore, two different cases with non-obstructive azoospermia manifested significant down-regulation of the germ cell marker gene expression. Also, comparative analysis between two ages of mice showed different patterns of expression of DDX4. These findings suggest that DDX4 as a male germ cell marker would be useful for future advanced investigations in reproductive medicine.

Conflict of Interest

The authors declare that they have no competing interests.

References

- Huang P, Wang T. Spermatogonial stem cell and TGF- β {eta} involved regulation of proliferation and differentiation. arXiv preprint arXiv:1706.03892. 2017 Jun 13.
- Yoshida S. Open niche regulation of mouse spermatogenic stem cells. *Dev Growth Differ*. 2018;60(9):542-52.
- Meinhardt A, Wang M, Schulz C, Bhushan S. Microenvironmental signals govern the cellular identity of testicular macrophages. *J leukoc Biol*. 2018;104(4):757-66.
- Wang F, Han D. Sertoli cell phagocytosis: an essential event for spermatogenesis. In: Wu W, Ziglioli F, Maestroni U, editors. *Male reproductive health*. UK: IntechOpen; 2019. p. 69-84.
- Chen H, Wang Y, Ge R, Zirkin BR. Leydig cell stem cells: Identification, proliferation and differentiation. *Mol Cell Endocrinol*. 2017;445:65-73.
- De Rooij DG. The nature and dynamics of spermatogonial stem cells. *Development*. 2017;144(17):3022-30.
- Ishikura Y, Yabuta Y, Ohta H, Hayashi K, Nakamura T, Okamoto I, et al. In vitro derivation and propagation of spermatogonial stem cell activity from mouse pluripotent stem cells. *Cell Rep*. 2016;17(10):2789-804.
- Zhou H, Zeng Z, Koentgen F, Khan M, Mombaerts P. The testicular soma of Tsc22d3 knockout mice supports spermatogenesis and germline transmission from spermatogonial stem cell lines upon transplantation. *Genesis*. 2019;57(6):e23295.

9. Chapman KM, Medrano GA, Jaichander P, Chaudhary J, Waits AE, Nobrega MA, et al. Targeted germline modifications in rats using CRISPR/Cas9 and spermatogonial stem cells. *Cell Rep.* 2015;10(11):1828-35.
10. Griswold MD. Spermatogenesis: the commitment to meiosis. *Physiol Rev.* 2016;96(1):1-17.
11. Robles V, Herraez P, Labbé C, Cabrita E, Pšenička M, Valcarce DG, et al. Molecular basis of spermatogenesis and sperm quality. *Gen Comp Endocrinol.* 2017;245:5-9.
12. Toocheck C, Clister T, Shupe J, Crum C, Ravindranathan P, Lee TK, et al. Mouse spermatogenesis requires classical and nonclassical testosterone signaling. *Biol Reprod.* 2016;94(1):11.
13. Silber S. Hormonal regulation of spermatogenesis. In: *Fundamentals of male infertility.* USA: Springer; 2018. p. 39-41.
14. Miller SR, Morales MN, Frost KL, Cherrington NJ, Wright SH. Cell model for studying nucleoside transporters, a key component of the blood-testis barrier. *FASEB J.* 2019;33(21):507-12.
15. Hui L, Nie Y, Li S, Guo M, Yang W, Huang R, et al. Matrix metalloproteinase 9 facilitates Zika virus invasion of the testis by modulating the integrity of the blood-testis barrier. *PLoS Pathog.* 2020;16(4):e1008509.
16. Nayernia K, Lee JH, Drusenheimer N, Nolte J, Wulf G, Dressel R, et al. Derivation of male germ cells from bone marrow stem cells. *Lab Invest.* 2006;86(7):654-63.
17. Nayernia K, Nolte J, Michelmann HW, Lee JH, Rathsack K, Drusenheimer N, et al. In vitro-differentiated embryonic stem cells give rise to male gametes that can generate offspring mice. *Dev Cell.* 2006;11(1):125-32.
18. Cardoso HJ, Figueira MI, Socorro S. The stem cell factor (SCF)/c-KIT signalling in testis and prostate cancer. *J Cell Commun Signal.* 2017;11(4):297-307.
19. Azizi H, Hamidabadi HG, Skutella T. Differential proliferation effects after short-term cultivation of mouse spermatogonial stem cells on different feeder layers. *Cell J.* 2019;21(2):186-93.
20. Li R, Vannitamby A, Yue SS, Handelsman D, Hutson J. Mouse minipuberty coincides with gonocyte transformation into spermatogonial stem cells: a model for human minipuberty. *Reprod Fertil Dev.* 2017;29(12):2430-6.
21. Azizi H, Asgari B, Skutella T. Pluripotency potential of embryonic stem cell-like cells derived from mouse testis. *Cell J.* 2019;21(3):281-9.
22. Patel H, Bhartiya D. Direct action of FSH on testicular stem cells. *Stem Cell Res Ther.* 2019;10(1):261.
23. Ng CL, Yue Q, Cordula S. Notch signaling is required for survival of the germline stem cell lineage in testes of drosophila melanogaster. *BioRxiv.* 2019 Jan 1:682773.
24. Boellaard WP, Stoop H, Gillis AJ, Oosterhuis JW, Looijenga LH. VASA mRNA (DDX4) detection is more specific than immunohistochemistry using poly-or monoclonal antibodies for germ cells in the male urogenital tract. *Medicine (Baltimore).* 2017;96(30):e7489.
25. Sharma S, Schlatt S, Van Pelt A, Neuhaus N. Characterization and population dynamics of germ cells in adult macaque testicular cultures. *PLoS One.* 2019;14(6):e0218194.
26. Abd Razak R, Isa LM, Razali RM, Yusuf AM, Abd Wahab AY, Ariffin SA, et al. Identification of VASA gene expression in in-vitro culture from non-obstructive azoospermia (NOA) testicular biopsy cells: a study to unlock knowledge of male infertility. *IIUM Med J Malaysia.* 2018;17(2).
27. Li M, Zhu F, Li Z, Hong N, Hong Y. Dazl is a critical player for primordial germ cell formation in medaka. *Sci Rep.* 2016;6(1):1-7.
28. Schwager EE, Meng Y, Extavour CG. vasa and piwi are required for mitotic integrity in early embryogenesis in the spider Parasteatoda tepidariorum. *Dev Biol.* 2015;402(2):276-90.
29. Tanaka SS, Toyooka Y, Akasu R, Katoh-Fukui Y, Nakahara Y, Suzuki R, et al. The mouse homolog of Drosophila Vasa is required for the development of male germ cells. *Genes Dev.* 2000;14(7):841-53.
30. Wang K, Zhang H, Hu Q, Shao C, Chen S. Expression and purification of half-smooth tongue sole (*Cynoglossus semilaevis*) CSDAZL protein. *Protein Expr Purif.* 2014;102:8-12.
31. Zhu W, Wang T, Zhao C, Wang D, Zhang X, Zhang H, et al. Evolutionary conservation and divergence of Vasa, Dazl and Nanos1 during embryogenesis and gametogenesis in dark sleeper (*Odontobutis potamophila*). *Gene.* 2018;672:21-33.
32. Lejong M, Choa-Duterte M, Vanmuylder N, Louryan S. Is Vasa such a highly specific marker for primordial germ cells? a comparison of VASA and HSP90 proteins expression in young chicken embryos. *Morphologie.* 2020;104(344):20-6.
33. Guo J, Cairns BR. Isolation and enrichment of spermatogonial stem cells from human testis tissues. *Curr Protoc Stem Cell Biol.* 2019;49(1):e77.

34. Murdock MH, David S, Swinehart IT, Reing JE, Tran K, Gassei K, et al. Human testis extracellular matrix enhances human spermatogonial stem cell survival in vitro. *Tissue Eng Part A*. 2019;25(7-8):663-76.
35. Mossadegh-Keller N, Sieweke MH. Characterization of mouse adult testicular macrophage populations by immunofluorescence imaging and flow cytometry. *Bio Protoc*. 2019;9(5):10.21769/Bio Protoc.3178.
36. Li X, Sun T, Wang X, Tang J, Liu Y. Restore natural fertility of Kit w/Kit wv mouse with nonobstructive azoospermia through gene editing on SSCs mediated by CRISPR-Cas9. *Stem Cell Res Ther*. 2019;10(1):271.
37. Abbasi H, Tahmoorespur M, Hosseini SM, Nasiri Z, Bahadorani M, Hajian M, et al. THY1 as a reliable marker for enrichment of undifferentiated spermatogonia in the goat. *Theriogenology*. 2013;80(8):923-32.
38. Castrillon DH, Quade BJ, Wang TY, Quigley C, Crum CP. The human VASA gene is specifically expressed in the germ cell lineage. *Proc Natl Acad Sci USA*. 2000;97(17):9585-90.
39. Lee WY, Lee R, Park HJ, Do JT, Park C, Kim JH, et al. Characterization of male germ cell markers in canine testis. *Anim Reprod Sci*. 2017;182:1-8.
40. Hashimoto H, Sudo T, Mikami Y, Otani M, Takano M, Tsuda H, et al. Germ cell specific protein VASA is over-expressed in epithelial ovarian cancer and disrupts DNA damage-induced G2 checkpoint. *Gynecol Oncol*. 2008;111(2):312-9.
41. Zhang T, Ji L, Liu B, Guan W, Liu Q, Gao Y. Testicular germ cell tumors: a clinicopathological and immunohistochemical analysis of 145 cases. *Int J Clin Exp Pathol*. 2018;11(9):4622-9.
42. Guo X, Gui YT, Tang AF, Lu LH, Gao X, Cai ZM. Differential expression of VASA gene in ejaculated spermatozoa from normozoospermic men and patients with oligozoospermia. *Asian J Androl*. 2007;9(3):339-44.
43. Ando M, Yamaguchi K, Chiba K, Miyake H, Fujisawa M. Expression of VASA mRNA in testis as a significant predictor of sperm recovery by microdissection testicular sperm extraction in patient with nonobstructive azoospermia. *J Androl*. 2012;33(4):711-6.