



An Incidental Detection of a Cryptic Complex Chromosome Rearrangement Found During NGS Based PGT-SR: A Case Report

Leyla Özer^{1,2*}, Suleyman Aktuna^{1,2}, Evrim Unsal^{1,2}, Aysun Baltaci³, Volkan Baltaci^{1,2}

1- Department of Medical Genetics, Faculty of Medicine, Yuksek İhtisas University, Ankara, Turkey

2- Mikrogen Genetic Diagnosis Center, Ankara, Turkey

3- Gen-Art IVF Center, Ankara, Turkey

Abstract

Background: Complex chromosome rearrangements (CCRs) involve more than 2 chromosomal breakpoints and cause the exchanges of chromosomal segments between two or more chromosomes. The carriers of CCRs have normal phenotypes, but they have a higher risk of reproductive failure.

Case Presentation: This paper presents a couple with a history of two affected children, one spontaneous abortion, three in vitro fertilization (IVF) failures, and one healthy boy who were referred to our laboratory for preimplantation genetic testing (PGT). The wife had been evaluated as a carrier of 46,XX,t(2;6)(p21;p25); therefore, four IVF treatment cycles supported with PGT for this translocation had been performed in different IVF centers until the couple consulted our laboratory. Only one of these four IVF attempts had resulted in a healthy boy and this IVF study had been performed with fluorescence in situ hybridization (FISH)-based preimplantation genetic testing for structural chromosomal rearrangements (PGT-SR). The fifth IVF study with next-generation sequencing (NGS)-based PGT was performed by our laboratory and no healthy embryo was found in evaluated 6 embryos. During our NGS-based PGT, the cryptic involvement of 12p was firstly detected. FISH with chromosome 2,6, and 12 specific probes revealed that the mother was a carrier of a balanced 3-way translocation of 46,XX,t(2;6;12)(p21;p25;p13).

Conclusion: NGS based PGT-SR method is an accurate method for detecting the copy number variations and is helpful to find out the cryptic CCRs.

Keywords: Chromosomal translocation, Chromosome abnormalities, Next generation sequencing, Preimplantation genetic testing.

To cite this article: Özer L, Aktuna S, Unsal E, Baltaci A, Baltaci V. An Incidental Detection of a Cryptic Complex Chromosome Rearrangement Found During NGS Based PGT-SR: A Case Report. *J Reprod Infertil.* 2022;23(4):303-309. <https://doi.org/10.18502/jri.v23i4.10817>.

Introduction

Chromosomal structural anomalies usually involve two or more chromosomes, and complex chromosome rearrangements (CCRs) indicate three or more chromosome breaks. CCRs are rarely seen and can be familial or de novo. CCRs are classified according to the number of chromosomes involved, the number of chromosome breaks, and the involvement of intrachromo-

somal or interchromosomal insertions. CCRs are usually classified as three-way rearrangements, double two-way translocations, and exceptional CCRs (1-3). Complex chromosomal translocations involve more than the reciprocal exchange of segments between two chromosomes. These can include double translocation, three-way translocation, and more complex rearrangements with

* Corresponding Author:
Leyla Özer, Department of
Medical Genetics, Faculty
of Medicine, Yuksek İhtisas
University, Ankara, Turkey
E-mail:
leyla_ozer@yahoo.com

Received: Dec. 25, 2021

Accepted: May 5, 2022

terminal exchange, inversion, interstitial insertion, deletion, and more than three chromosomes. Carriers of balanced CCRs have a high risk of recurrent pregnancy losses or affected children. Balanced CCR carriers are phenotypically normal; in fact, a low probability of having a child with a normal balanced karyotype was reported. Several studies (4, 5) found a 50% chance of spontaneous miscarriage and a 20% chance of having chromosomally abnormal children. A wide variety of possible gametes are expected due to the number of chromosomes and the number of breakpoints involved. Hu et al. postulated that the odds of balanced or normal embryos in couples with balanced CCRs are very low (less than 6%) (6). Scriven et al. reported that female CCR carriers have a lower proportion of normal/balanced embryos (4.9%) compared with male carriers (19.4%). Preimplantation genetic testing (PGT) is a good option for these CCR carriers to select normal/balanced embryos (7).

Preimplantation genetic testing is performed to analyze embryos for monogenic disorders and chromosomal abnormalities before embryo transfer. PGT is a group of genetic tests that are divided into three categories: PGT for aneuploidy (PGT-A), PGT for monogenic/single gene defects (PGT-M), and PGT for chromosomal structural rearrangements (PGT-SR) (7-9). PGT-SR is used to screen unbalanced embryos for balanced structural chromosomal abnormality carriers (translocation, inversion). The genetic techniques used for PGT are fluorescence in-situ hybridization FISH, array comparative genomic hybridization (array CGH), and next-generation sequencing (NGS) methods. Recently, array CGH and NGS have been commonly used due to their advantage in checking all 24 chromosomes for PGT. NGS and array CGH detect aneuploidies and unbalanced chromosomal structural abnormalities (copy number variants) simultaneously (9, 10).

In this paper, a PGT-SR study of a couple was reported where the wife was a carrier of a balanced translocation. During the NGS study of embryos, it was found that she was a carrier of a 3-way complex translocation and in this translocation, the cryptic involvement of 12p13.33p13.31 was observed.

Case Presentation

The nonconsanguineous couple was referred from the Genart IVF Center (Ankara, Turkey) to our genetic laboratory (Mikrogen Genetic Diag-

nosis Center, Ankara, Turkey) in 2020. The couple with a history of two affected children, one spontaneous abortion, three in vitro fertilization (IVF) failures, and one healthy boy was referred to our laboratory for preimplantation genetic testing (PGT) (Figure 1). The wife had been evaluated as a carrier of 46, XX, t (2; 6) (p21; p25); therefore, four IVF treatment cycles supported by PGT for this translocation had been performed in different IVF centers until the couple consulted our laboratory. The first child of the family was an affected girl who suffered from epilepsy and spasticity and died at 7 months old. The second pregnancy resulted in spontaneous abortion. The second child was a girl with bilateral anophthalmia, epilepsy, and multiple congenital anomalies who died at 3.5 years old. After the loss of her second child, it was noticed that the mother was a carrier of the 46,XX,t (2;6) (p21; p25) translocation. Since the mother was found to be a carrier of the translocation, four IVF treatment cycles supported by PGT-SR had been performed in different IVF centers until the couple consulted our laboratory. In their first IVF study, seven embryos had been detected by FISH analysis (using 2p telomeric, chromosome 2 centromeric, and chromosome 6 centromeric probes) and only one healthy embryo had been found. Trisomy 2p in three embryos, monosomy 2p in one embryo, trisomy 2 in one embryo, and monosomy 2 in one embryo have been reported. The family got a healthy boy from this first IVF treatment with FISH-based PGT-SR and their son is 9 years old now. In the second IVF attempt with the NGS-based PGT-SR, 4 embryos had been detected and no healthy embryos had been found (monosomy 6 in embryo 1, monosomy 2 in embryo 2, monosomy 11 in embryo 3 and monosomy 2p, trisomy 6p, and trisomy 12 in embryo 4). In the third IVF treatment with NGS-based PGT-SR, 4 embryos had been detected and no healthy embryos had been found (monosomy 2 in embryo 1, monosomy 2p, trisomy 6p in embryo 2, trisomy 6p in embryo 3,




Figure 1. The pedigree analysis of the family

and monosomy 2p, trisomy 16 in embryo 4). In the fourth IVF treatment with the NGS-based PGT-SR, four embryos were evaluated but no healthy embryos were found. Since the report of this study did not reach us, detailed results could not be outlined. The couple was referred to our laboratory to perform NGS-based PGT-SR as their 5th IVF treatment.

PGT-SR study: Ovarian stimulation was performed using standard techniques of gonadotrophin-releasing hormone (GnRH) agonist down-regulation combined with ovarian stimulation using a combination of recombinant FSH (Gonal-f; EMD Serono Inc., Germany) and urinary-derived gonadotrophin stimulation (Menapur; Ferring Pharmaceuticals, Germany; Merioval; IBSA Institut Biochimique SA, Switzerland). Oocyte retrieval occurred approximately 36 hr after human chorionic gonadotrophin (HCG) administration and was performed via ultrasound-guided transvaginal aspiration. Spermatozoa used for the IVF procedure were prepared using standard density gradient centrifugation (Isolate; Irvine Scientific, USA).

In order to prepare for the PGT study, 4-5 trophoblast cells were biopsied from blastocyst-stage embryos on day 5. The biopsied cells were lysed and the whole genome amplification (WGA) was performed with the Ion ReproSeq PGS kit (Thermo Scientific, USA) including fragmentation, library preparation, and amplification. All six biopsied blastocysts were successfully amplified. Next generation sequencing was performed on the Ion GeneStudio S5 (Thermo Scientific, USA) platform according to the manufacturer's protocol. Results were analyzed with Ion Reporter Software. FISH analysis was carried out on metaphase and interphase nuclei of the mother obtained from PHA stimulated lymphocyte cultures by using commercial FISH probes for 2p telomeric (green), 2q telomeric (orange), 6p telomeric (green), 6 centromeric aqua and 12p telomeric (green), and 12q telomeric (orange) probes.

Results

PGT-SR results: NGS analysis was done for 6 embryos and no healthy embryo was found. The results of this study were:

Embryo 1; 47,XX,del(12)(p13.33p13.31),+15
Embryo 2; 47,XX,del(6)(p25.3p25.2),dup(12)(p13.33p13.31),+15

Embryo 3; 45,XY, dup(2)(p25.3p21), del(6)(p25.3p25.2),-22

Embryo 4; 47,XX, dup(2)(p25.3p21), del(6)(p25.3p25.2),+16

Embryo 5; chaotic (multiple aneuploidies including chromosome 1, 2, 5, 6, 9, 12, 15, 19)

Embryo 6; chaotic (multiple aneuploidies including chromosome 2, 6, 7, 8, 9, 13, 16, 17, 19, 20, 21, 22) (Figure 2).

As a result of NGS analysis, cryptic involvement of chromosome 12p13.33p13.31 region was detected. The breakpoints and sizes of the variants are listed below:

chr12 (p13.33p13.31): 145,739-6,189,570 approximately 6MB

chr6 (p25.3p25.2): 60,000-4,061,907 approximately 4MB

chr2 (p25.3p21): 10,000-45,356,328 approximately 45 MB

Diagnosis of the chromosomes involved in complex chromosomal rearrangements: After PGT-SR analysis, the karyotype and FISH analysis of the mother was studied to exclude possible complex translocation. FISH analysis with chromosome 2, 6, and 12 specific probes revealed that the mother was a carrier of a balanced 3-way translocation of 46,XX,t (2;6;12)(p21;p25;p13). The cryptic involvement of 12p13 was first detected during our PGT-based NGS study of embryos. G-banded metaphases were captured (Figure 3), and a subsequent metaphase FISH study was prepared. The results showed the presence of a 12p green signal on derivative chromosome 2 and a 12q red signal on chromosome 12 (Figure 4A). Subsequent FISH on the other metaphases (Figure 4B) showed the presence of a 2p green signal on derivative chromosome 6 and a 2q red signal on chromosome 2. Figure 4C represents the presence of a 6p green signal on the short arm of chromosome 12. This confirmed the complex cryptic translocation of 46,XX,t(2;6;12)(p21;p25;p13), which was not detected on karyotyping and was incidentally found during the PGT-SR NGS study. The karyotype of the mother after metaphase FISH was therefore 46,XX,t(2;6;12) (p21;p25;p13).

Discussion

Balanced complex translocation carriers of three-way exchange involving three chromosomes increase the risk of recurrent miscarriages and affect children due to the production of unbalanced gametes (11).




Figure 2. PGT-SR results: A) positive for heterozygous ~6 Mb deletion at 12 p13.33p13.31, trisomy 15; B) positive for heterozygous ~4 Mb deletion at 6 p25.3p25.2, positive for heterozygous ~4 Mb duplication at 12 p13.33p13.31, trisomy 15; C) positive for heterozygous ~4 Mb deletion at 6 p25.3p25.2, positive for heterozygous ~45 Mb duplication at 2p25.3p21, monosomy; 22 D) positive for heterozygous ~4 Mb deletion at 6 p25.3p25.2, positive for heterozygous ~45 Mb duplication at 2p25.3p21, trisomy; 16 E) chaotic (multiple aneuploidies); and F) chaotic (multiple aneuploidies)

The estimated frequency of balanced complex chromosomal translocations involving three chromosomes was reported as 0.1% among couples with a history of pregnancy losses (14). Li et al. searched for the influence of three types of balanced CCR on early embryonic development and molecular karyotype, and they found that the frequency of high quality blastocysts for three-way rearrangements was low (15). Scriven et al. reported that the chance of normal/balanced embryos for male and female three-way translocation carriers was 19.4% and 4.9%, respectively (7). In our case, the carrier mother had a history of af-

ected children, miscarriages, and IVF failure. CCR carriers have an increased risk of chromosomal instability in embryos, so identification of the chromosomes involved in the translocation is important in providing the CCR couple with accurate genetic counselling for reproductive risks and PGT options. The couple's chance of having a normal or balanced embryo is expected to be very low. When the couple was referred to our laboratory, four PGT-SR studies had previously been performed in other laboratories due to the wife's two-way translocation. One of them was the FISH based PGT-SR study, which only detects the im-




Figure 3. A) karyotype analysis of the mother, GTG-banded chromosomal analysis reveals the presence of a balanced translocation involving chromosome 2, 6, and 12; B) the exchanged segments were represented in ideogram

balances due to two-way translocation between chromosomes 2 and 6; therefore, the results of the current study cannot yield information about the participation of chromosome 12. Among the three NGS based PGT-SR studies, none of them could find a chromosomal abnormality involving the




Figure 4. FISH images: A) Metaphase FISH showing 12p (green) signal on 2p (green signal: 12p, red signal: 12q), B) Metaphase FISH showing 2p (green) signal on 6p (green signal: 12p, red signal: 2q), C) Metaphase FISH showing 6p (green) signal on 12p (green signal: 6p, red signal: 12q, aqua signal: 6 centromere)

12p13.33p13.31 region. Trisomy 12 was only reported in one of them. Comparing the results of our PGT-SR study with previous NGS-based PGT-SR studies in couples, it was found that copy number variations of 12p13.33p13.31 were not

reported before the current study. CCR was suspected as a result of our NGS-based PGT study, and after the PGT-SR study, the presence of three-way translocation of the wife was confirmed by FISH analysis. If the presence of the complex chromosomal translocation had been detected in previous IVF applications, the family could have been better informed about their reproductive risks.

Some of the CCRs have cryptic changes like unexpected involvement of chromosomes or micro-deletions or microduplications. Several studies (1, 11, 12) found that seemingly balanced reciprocal translocations can have unexpected complexity and cryptic changes. The accuracy for detecting chromosomes and chromosomal breakpoints depends on the quality of the chromosome analysis. Conventional cytogenetic analysis may have some difficulties in detecting complex chromosome rearrangements because of unexpected involvement of chromosomes and submicroscopic chromosomal anomalies. Telomeric regions of all chromosomes have the same light-staining pattern of G-banding, so conventional cytogenetics may miss these small translocations, which are called "cryptic translocations". Conventional cytogenetics is inadequate for detecting these translocations since the size and banding pattern of the chromosome regions involved are too similar for the exchange to be detected, or the size of the exchanged segments is smaller than the resolution limit of the G-banding. With the use of FISH and array technologies, it is possible to define complex chromosomal translocations (11). In our case, cryptic involvement of the 12p13.33p13.31 segment was first suspected according to the NGS based PGT-SR study's results and then confirmed by FISH in lymphocyte cultured cells of the wife.

The reason why the involvement of chromosome 12 was overlooked during previous PGT studies was thought to be related to the size and banding pattern of the chromosome 6 and chromosome 12 segments involved in the translocation. The resolution in conventional cytogenetic analysis is approximately 5–10 MB according to banding quality. The translocated segments of chromosomes 6 and 12 are approximately the same size and have the same banding pattern (light band), thus making it difficult to recognize the presence of complex translocation. As the CCR carrier status of the wife was not known before our PGT-SR study, the NGS-based PGT-SR study helped us to find the presence of CCR. Previous studies show-

ed that cryptic chromosome involvement in some CCRs carriers was diagnosed during PGT workup incidentally (16, 20). PGT-SR for cryptic translocations detects a higher frequency of unbalanced karyotypes in comparison with prenatal diagnosis techniques (chorionic villus sampling and amniocentesis) (13).

Although it is difficult to find normal embryos for balanced complex chromosome rearrangement (BCCR) carriers, preimplantation genetic testing gives a chance for carriers of CCR to reduce the risk of recurrent miscarriages and having offspring with unbalanced chromosomal changes (12). A limited number of cases of PGT for carriers of CCRs were reported. Some of the cases were analyzed by FISH technique (16, 17), but with the development of NGS technologies, the number of PGD cases analyzed by NGS is increasing (15, 18-20). FISH, array CGH, and NGS methods are commonly used for PGT-SR studies. FISH, as compared with NGS, has the disadvantage of screening only a limited number of chromosomes due to the limited number of probes for simultaneous use. PGT-SR using NGS technology detects not only chromosomes that participate in translocation but also recognizes all involved chromosomes. With the development of molecular technology, NGS became an accurate method for detecting the copy number variations in embryos.

Conclusion

In our reported case, balanced three-way translocation carrier status was detected after NGS-based PGT-SR, in spite of having two-way balanced translocation. The sizes and banding patterns of exchanged fragments of chromosome 6 and 12 probably caused some difficulty in defining the correct diagnosis. In conclusion, our study showed that the NGS-based PGT technique can effectively help families with a cryptic balanced translocation plan their pregnancies and detect unbalanced chromosomal abnormalities that may be below the resolution limit of conventional cytogenetic analysis. FISH and conventional karyotype analyses are the gold standards for detecting balanced CCR carriers. Although the FISH technique is still accepted as the gold standard method for PGT-SR study of CCRs, the NGS-based PGT-SR method provides the opportunity to find the cryptic CCRs incidentally; therefore, this case report also emphasizes the importance of simultaneous scanning of all chromosomes to detect cryptic

CCRs. The use of the FISH and NGS methods together in the preimplantation genetic diagnosis process will help to determine the potential risks and make a more accurate diagnosis. The detection of undiagnosed complex balanced translocation carriers among IVF patients will likely be improved by NGS-based PGT-ST.

Conflict of Interest

Authors declare no conflict of interest.

References

- Pellestor F, Anahory T, Lefort G, Puechberty J, Liehr T, He'don B, et al. Complex chromosomal rearrangements: origin and meiotic behavior. *Hum Reprod Update*. 2011;17(4):476-94.
- Madan K. Balanced complex chromosome rearrangements: reproductive aspects. a review. *Am J Med Genet Part A*. 2012;158A(4):947-63.
- Madan K. What is a complex chromosome rearrangement? *Am J Med Genet A*. 2013;161A(5):1181-4.
- Gorski JL, Kistenmacher ML, Punnett HH, Zackai EH, Emanuel BS. Reproductive risks for carriers of complex chromosome rearrangements: analysis of 25 families. *Am J Med Genet*. 1988;29(2):247-61.
- Madan K, Nieuwint AW, Van Bever Y. Recombination in a balanced complex translocation of a mother leading to a balanced reciprocal translocation in the child. Review of 60 Cases of balanced complex translocations. *Hum Genet*. 1997;99(6):806-15.
- Hu L, Wei Y, Luo K, Xie P, Gong F, Xiong B, et al. Clinical outcomes in carriers of complex chromosomal rearrangements: a retrospective analysis of comprehensive chromosome screening results in seven cases. *Fertil Steril*. 2018;109(3):486-92.
- Scriven PN, Bint SM, Davies AF, Ogilvie CM. Meiotic outcomes of three-way translocations ascertained in cleavage-stage embryos: refinement of reproductive risks and implications for PGD. *Eur J Hum Genet*. 2014;22(6):748-53.
- Harton G, Braude P, Lashwood A, Schmutzler A, Traeger-Synodinos J, Wilton L, et al. ESHRE PGD consortium best practice guidelines for organization of a PGD centre for PGD/preimplantation genetic screening. *Hum Reprod*. 2010;26(1):14-24.
- ESHRE PGT consortium steering committee, Carvalho F, Coonen E, Goossens V, Kokkali G, Rubio C, et al. ESHRE PGT consortium good practice recommendations for the organisation of PGT. *Hum Reprod Open*. 2020;2020(3):hoaa021.
- [No Authors listed]. Preimplantation genetic testing: ACOG committee opinion, Number 799. *Obstet Gynecol*. 2020;135(3):e133-e7.
- Gregori MD, Ciccone R, Magini P, Pramparo T, Gimelli S, Messa J, et al. Cryptic deletions are a common finding in "balanced" reciprocal and complex chromosome rearrangements: A study of 59 patients. *J Med Genet*. 2007;44(12):750-62.
- Escudero T, Estop A, Fischer J, Munne S. Preimplantation genetic diagnosis for complex chromosome rearrangements. *Am J Med Genet A*. 2008;146A(13):1662-9.
- Vanneste E, Melotte C, Voet T, Robberecht C, Debrock S, Pexsters A, et al. PGD for a complex chromosomal rearrangement by array comparative genomic hybridization. *Hum Reprod*. 2011;26(4):941-9.
- Nonaka T, Ooki I, Enomoto T, Takakuwa K. Complex chromosomal rearrangements in couples affected by recurrent spontaneous abortion. *Int J Gynaecol Obstet*. 2015;128(1):36-9.
- Li G, Shi W, Niu W, Xu J, Guo Y, Su Y, et al. The influence of balanced complex chromosomal rearrangements on preimplantation embryonic development potential and molecular karyotype. *BMC Genomics*. 2020;21(1):326.
- Athalye A, Sanap R, Madon P, Naik D, Warang D, Padyal P, et al. Preimplantation genetic testing for a complex chromosome rearrangement, case report of a cryptic translocation detected on pre-PGT workup. *Obstet Gynecol Int J*. 2018;9(2):138-41.
- McKenzie LJ, Cisneros PL, Torsky S, Bacino CA, Buster JE, Carson SA, et al. Preimplantation genetic diagnosis for a known cryptic translocation: follow-Up clinical report and implication of segregation products. *Am J Med Genet*. 2003;121A(1):56-9.
- Brunet B, Shen J, Cai L, Xie J, Cui Y, Liu J, et al. Preimplantation genetic testing for complex chromosomal rearrangement carriers by next-generation sequencing. *Reprod Biomed Online*. 2018;37(3):375-82.
- Wang X, Wu C, Hao D, Zhang J, Tan C, Cheng D, et al. One healthy live birth after preimplantation genetic testing of a cryptic balanced translocation (9;13) in a family with cerebral palsy and glaucoma: a case report. *BMC Med Genomics*. 2021;14(1):82.
- Frumkin T, Peleg S, Gold V, Reches A, Asaf S, Azem F, et al. Complex chromosomal rearrangement-a lesson learned from PGS. *J Assist Reprod Genet*. 2017;34(8):1095-1100.