

Exposure to Environmental Organic Mercury and Impairments in Human Fertility

Geir Bjørklund^{1*}, Jan Aaseth^{2,3}, Maryam Dadar⁴, Monica Butnariu^{5,6}, Salvatore Chirumbolo⁷

1- Council for Nutritional and Environmental Medicine, Mo i Rana, Norway

2- Research Department, Innlandet Hospital Trust, Brumunddal, Norway

3- Inland Norway University of Applied Sciences, Elverum, Norway

4- Razi Vaccine and Serum Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Karaj, Iran

5- Banat's University of Agricultural Sciences and Veterinary Medicine "King Michael I of Romania" from Timisoara, Timisoara, Romania

6- CONEM Romania Biotechnology and Environmental Sciences Group, Banat's University of Agricultural Sciences and Veterinary Medicine "King Michael I of Romania" from Timisoara, Timisoara, Romania

7- Department of Neurological and Movement Sciences, University of Verona, Verona, Italy

* Corresponding Author: Geir Bjørklund, Council for Nutritional and Environmental Medicine, Toften 24, 8610 Mo i Rana, Norway
E-mail: bjorklund@conem.org

To cite this article: Bjørklund G, Aaseth J, Dadar M, Butnariu M, Chirumbolo S. Exposure to Environmental Organic Mercury and Impairments in Human Fertility. *J Reprod Infertil.* 2019;20(3):195-197.

It is widely known that mercury (Hg) is highly toxic to humans. Environmental Hg pollution still represents a huge health concern worldwide (1). Human industrial activities have led to raised levels of Hg in the air, soil, and fresh and sea waters, and to bioaccumulation along the food chain. Humans easily absorb Hg through fish consumption. Also, it is a major toxicant because it has adverse effects on human reproductive health (2). Mercury, particularly in its organic forms, methyl-Hg and ethyl-Hg, is toxic even when individuals are exposed to relatively low Hg levels (3). Toxic effects of Hg can easily be observed in the nervous, digestive, and immune systems, in addition to lungs, kidneys, skin, and eyes, as reported by the World Health Organization (4). Many of these effects are mediated by the host's immune response via its limited detoxification and metal excretion capabilities. Genetic polymorphism of scavenging Hg-complexed proteins are factors that can induce stressor effects, which also depend on the capability of enzymatic detoxification systems, aside from the dose of the toxicant (5).

The evidence that organic Hg easily crosses the placental barrier and reaches the fetus represents another great concern. Thereby, Hg is a major cause of neural developmental defects including delayed postnatal development (6). Both elemental Hg⁰ and methyl-Hg easily cross the placenta, and it should be emphasized that the developing fetal brain is the most sensitive organ. However,

the more recent debate about the toxicity of dietary Hg in fetal or prenatal toxicology (7) has revealed controversies about Hg effects on fetal development, particularly because it is difficult to establish a reliable Hg-pharmacokinetics in adult women undergoing pregnancy. The pharmacokinetics and threshold of plasma Hg-acceptable levels in pregnancy are still matters of debate, and updated evidence-based official guidelines are generally lacking (8).

Recent reports together with the previous information from the Minimata and Iraq disasters have reignited this debate (9). Mercury exposure may explain cases of idiopathic infertility in males as well as in females (10). Moreover, it affects chromosomes or embryonic cell DNA, presumably by its interference with one-carbon transfer since a methionine carrier transports it across membranes (11). Recent evidence has reported that Hg inhibits the energy metabolism of spermatozoa, thus preventing their normal functioning (12). Methyl-Hg causes reproductive organ damage in women (13).

Furthermore, menstrual disorders, sterility, and spontaneous abortion following abnormal Hg toxicology have been reported (14). Spontaneous abortion can occur when the pregnant woman is exposed to organic Hg or other toxic agents (15). Many spontaneous abortions do not have a known cause, but reports indicate that Hg exposure might be a triggering factor in several cases (16). Paternal exposure to Hg, and to other industrial chemi-

cals, and pesticides are also presumed to play a role 17). Studies on a Chinese population revealed that neural tube defects were associated with exposure to methyl-Hg in pregnancy and higher placental deposition of the compound (18). Based on the current knowledge, great concerns have been raised regarding Romanian districts with remarkably high Hg pollution (19).

Regarding pre-and postnatal risks of Hg exposure, it is relevant that the concentrations of Hg in the nervous system and other target organs of children may be higher than in the mothers, in part due to the small weight of a newborn child associated with a high rate of gastrointestinal absorption and low renal excretion, implying that hazardous Hg exposure may continue after birth (20).

Taken together, the comments presented here illustrate that the relationship between Hg toxicology and fertility has still been scarcely addressed in the literature, despite previous lessons from Minimata and some few reports in recent years. Apparently, Hg exposure may have a considerable impact on multiple stages of reproduction, from before conception to the maturation of organs and endocrine systems and further to the healthy development of the child. Therefore, a continued systematic investigation and assessment of the current state of knowledge about congenital and developmental anomalies with possible associations to environmental pollutants are of particular importance.

Conflict of Interest

The authors declare no conflict of interest.

References

1. Rana MN, Tangpong J, Rahman MM. Toxicodynamics of Lead, Cadmium, Mercury and Arsenic-induced kidney toxicity and treatment strategy: a mini review. *Toxicol Rep.* 2018;5:704-13.
2. Rahmani AM, Gia TN, Negash B, Anzanpour A, Azimi I, Jiang M, et al. Exploiting smart e-health gateways at the edge of healthcare Internet-of-things: a fog computing approach. *Future Gener Comput Syst.* 2018;78(2):641-58.
3. Bjørklund G, Dadar M, Mutter J, Aaseth J. The toxicology of mercury: current research and emerging trends. *Environ Res.* 2017;159:545-54.
4. World Health Organization. Mercury and health [Internet]. Geneva: World Health Organization; 2017 Mar 31. Available from: <https://www.who.int/news-room/fact-sheets/detail/mercury-and-health>.
5. Llop S, Ballester F, Broberg K. Effect of gene-mercury interactions on mercury toxicokinetics and neurotoxicity. *Curr Environ Health Rep.* 2015;2(2): 179-94.
6. Carocci A, Rovito N, Sinicropi MS, Genchi G. Mercury toxicity and neurodegenerative effects. *Rev Environ Contam Toxicol.* 2014;229:1-18.
7. Taylor CM, Emond AM, Lingam R, Golding J. Prenatal lead, cadmium and mercury exposure and associations with motor skills at age 7 years in a UK observational birth cohort. *Environ Int.* 2018;117: 40-7.
8. Golding J, Gregory S, Iles-Caven Y, Hibbeln J, Emond A, Taylor CM. Associations between prenatal mercury exposure and early child development in the ALSPAC study. *Neurotoxicology.* 2016;53:215-22.
9. Syversen T, Kaur P. The toxicology of mercury and its compounds. *J Trace Elem Med Biol.* 2012;26(4): 215-26.
10. Choy CM, Lam CW, Cheung LT, Briton-Jones CM, Cheung LP, Haines CJ. Infertility, blood mercury concentrations and dietary seafood consumption: a case-control study. *BJOG.* 2002;109 (10):1121-5.
11. Simmons-Willis TA, Koh AS, Clarjson TW, Ballatori N. Transport of a neurotoxicant by molecular mimicry: the methylmercury-L-cysteine complex is a substrate for human L-type large neutral amino acid transporter (LAT) 1 and LAT2. *Biochem J.* 2002;367(Pt 1):239-46.
12. Altunkaynak ME, Akgül N, Yahyazadeh A, Altunkaynak BZ, Türkmen AP, Akgül HM, et al. A stereological and histopathological study of the effects of exposure of male rat testes to mercury vapor. *Biotech Histochem.* 2015;90(7):529-34.
13. Wright DL, Afeiche MC, Ehrlich S, Smith K, Williams PL, Chavarro JE, et al. Hair mercury concentrations and in vitro fertilization (IVF) outcomes among women from a fertility clinic. *Reprod Toxicol.* 2015;51:125-32.
14. García-Forte P, Cohen-Corcía I, Córdoba-Doña JA, Reche-Rosado A, González-Mesa E. Toxic elements in hair and in vitro fertilization outcomes: a prospective cohort study. *Reprod Toxicol.* 2018; 77:43-52.
15. Yu E, Sim C, Park D, Koh Y, Heo J, Choe S, et al. Relationship between heavy metal concentration and number of spontaneous abortion experiences in Korean women: a retrospective study of the 6th Korean national health and nutrition examination survey. *Fertil Steril.* 2017;108(3):e320.
16. Anttila A, Sallmén M. Effects of parental occupational exposure to lead and other metals on spon-

- taneous abortion. *J Occup Environ Med.* 1995;37(8):915-21.
17. Mínguez-Alarcón L, Afeiche MC, Williams PL, Arvizu M, Tanrikut C, Amarasiwardena CJ, et al. Hair mercury (Hg) levels, fish consumption and semen parameters among men attending a fertility center. *Int J Hyg Environ Health.* 2018;221(2):174-82.
 18. Jin L, Liu M, Zhang L, Li Z, Yu J, Liu J, et al. Exposure of methyl mercury in utero and the risk of neural tube defects in a Chinese population. *Reprod Toxicol.* 2016;61:131-5.
 19. Neamtiu IA, Al-Abed SR, McKernan JL, Baciuc CL, Gurzau ES, Pogacean AO, et al. Metal contamination in environmental media in residential areas around Romanian mining sites. *Rev Environ Health.* 2017;32(1-2):215-20.
 20. Berlin M, Zalups RK, Fowler BA. Mercury. In: Nordberg GF, Fowler BA, Nordberg M, editors. *Handbook on the toxicology of metals.* Amsterdam: Elsevier; 2015. p.1013-69.